

# Engineered Cpf1 variants with altered PAM specificities

Linyi Gao<sup>1,2</sup> , David B T Cox<sup>1,3,4</sup>, Winston X Yan<sup>1,4,5</sup>, John C Manteiga<sup>3</sup>, Martin W Schneider<sup>1</sup> , Takashi Yamano<sup>6</sup>, Hiroshi Nishimasu<sup>6,7</sup>, Osamu Nureki<sup>6</sup>, Nicola Crosetto<sup>8</sup> & Feng Zhang<sup>1,2,9,10</sup> 

The RNA-guided endonuclease Cpf1 is a promising tool for genome editing in eukaryotic cells<sup>1–7</sup>. However, the utility of the commonly used *Acidaminococcus sp. BV3L6* Cpf1 (AsCpf1) and *Lachnospiraceae bacterium ND2006* Cpf1 (LbCpf1) is limited by their requirement of a TTTV protospacer adjacent motif (PAM) in the DNA substrate. To address this limitation, we performed a structure-guided mutagenesis screen to increase the targeting range of Cpf1. We engineered two AsCpf1 variants carrying the mutations S542R/K607R and S542R/K548V/N552R, which recognize TYCV and TATV PAMs, respectively, with enhanced activities *in vitro* and in human cells. Genome-wide assessment of off-target activity using BLISS<sup>7</sup> indicated that these variants retain high DNA-targeting specificity, which we further improved by introducing an additional non-PAM-interacting mutation. Introducing the identified PAM-interacting mutations at their corresponding positions in LbCpf1 similarly altered its PAM specificity. Together, these variants increase the targeting range of Cpf1 by approximately threefold in human coding sequences to one cleavage site per ~11 bp.

Programmable endonucleases from class 2 microbial CRISPR–Cas systems have enabled a wide range of applications in eukaryotic genome editing<sup>1–7</sup>. Recent work has demonstrated that in addition to the widely used type II-A Cas9, the type V-A system Cpf1 can mediate efficient genome editing. Cpf1 has several advantages compared to Cas9; for instance, it has low mismatch tolerance<sup>4–7</sup>, does not require a *trans*-activating crRNA, and can process its own CRISPR RNA (crRNA) array into mature crRNAs to facilitate targeting of multiple genes concurrently<sup>2,3</sup>.

We previously identified two orthologs of Cpf1 with robust activity in mammalian cells, *Acidaminococcus sp. BV3L6* Cpf1 (AsCpf1) and *Lachnospiraceae bacterium ND2006* Cpf1 (LbCpf1)<sup>1</sup>, both of which require a TTTV protospacer-adjacent motif (PAM), where V can be A, C, or G. For applications for which the location of the target site is critical, such as homology-directed repair or generation of loss-of-function mutations at specific exonic positions, the requirement

of a TTTV PAM may limit the availability of suitable target sites, reducing the practical utility of Cpf1. To address this limitation, we aimed to engineer variants of Cpf1 that can recognize alternative PAM sequences in order to increase its targeting range.


Previous work has shown that the PAM preference of Cas9 can be altered by mutations to residues in close proximity to the PAM DNA duplex<sup>8–11</sup>. We sought to investigate whether the PAM preference of Cpf1, despite its strong evolutionary conservation across different orthologs<sup>1</sup>, can also be modified. Based on the crystal structure of AsCpf1 in complex with crRNA and target DNA<sup>12</sup>, we selected 60 residues in AsCpf1 in proximity to the PAM duplex for targeted mutagenesis (Fig. 1a and Supplementary Table 1a). By randomizing the codons at each position using cassette mutagenesis, we constructed a plasmid library of AsCpf1 variants encoding most single amino acid substitutions at these residues. The use of codon randomization allowed us to attain greater mutational coverage than would have been expected with error-prone PCR, since it prevents representational bias caused by the template sequence.

To identify variants within this library with cleavage activity at non-canonical PAMs, we adapted a plasmid interference-based depletion screen in *Escherichia coli*<sup>1,8,13,14</sup> (Fig. 1b). In our modified assay, a pool of *E. coli*, with each bacterium expressing crRNA and a variant of Cpf1 from a plasmid maintained with chloramphenicol, was transformed with a second plasmid carrying an ampicillin-resistance gene and a target site bearing a mutated PAM. Successful cleavage of the second plasmid resulted in the loss of ampicillin resistance and subsequent cell death when grown on ampicillin-selective media. By comparing the sequences of the original library to the sequences of Cpf1-carrying plasmid DNA in surviving bacteria, we determined the variants that were depleted as a result of their novel cleavage activity of the mutated PAM.

To effectively use this approach to distinguish variants with non-canonical PAM activity from wild-type (WT) AsCpf1, we first determined PAM sequences at which WT AsCpf1 had minimal activity. We evaluated the tolerance of WT AsCpf1 to substitution mutations in the PAM, as determined by *E. coli* death due to successful plasmid interference. We focused on PAMs with single-nucleotide substitutions (i.e., NTTV, TNTV, and TTNV, where V was arbitrarily chosen to be C).

<sup>1</sup>Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA. <sup>2</sup>Department of Biological Engineering, Massachusetts Institute of Technology Cambridge, Massachusetts, USA. <sup>3</sup>Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA. <sup>4</sup>Harvard-MIT Division of Health Sciences and Technology, Harvard Medical School, Boston, Massachusetts, USA. <sup>5</sup>Graduate Program in Biophysics, Harvard Medical School, Boston, Massachusetts, USA. <sup>6</sup>Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, Japan. <sup>7</sup>JST, PRESTO, Tokyo, Japan. <sup>8</sup>Science for Life Laboratory, Division of Translational Medicine and Chemical Biology, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden. <sup>9</sup>McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA. <sup>10</sup>Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA. Correspondence should be addressed to F.Z. (zhang@broadinstitute.org).

Received 27 November 2016; accepted 16 May 2017; published online 5 June 2017; doi:10.1038/nbt.3900



**Figure 1** A bacterial interference-based negative selection screen identifies amino acid substitutions of AsCpf1 conferring activity at non-canonical PAMs. **(a)** Crystal structure of AsCpf1 (PDB ID: 5B43) in complex with crRNA and target DNA, highlighting the PAM nucleotides (magenta), and PAM-proximal residues selected for mutagenesis (blue). **(b)** Schematic of bacterial interference assay used to identify variants with altered PAM specificity. **(c)** Sensitivity of wild-type AsCpf1 to substitution mutations in the PAM as measured by bacterial interference. Error bars: mean  $\pm$  s.e.m. of  $n = 3$  plated transformations. **(d)** Scatter plots of screen readout, highlighting depleted variants. Each dot represents a distinct wild-type or mutant codon. The dashed lines indicate 15-fold depletion.

When transformed with NTTC and TCTC PAMs, *E. coli* expressing WT AsCpf1 had negligible survival on ampicillin media (**Fig. 1c**), indicating that these PAM sequences supported AsCpf1-mediated DNA plasmid cleavage and were not usable for screening the variant library. By contrast, the other five PAMs with a single mutation (TATC, TGTC, TTAC, TTCC, and TTGC) had notable survival rates. We subsequently screened the variant library for activity at these five PAMs, as well as an additional PAM with a double mutation (TCCC) (**Fig. 1d**).

Following deep sequencing readout, ~86% of the possible variants at the targeted residue positions were represented with at least 15 reads in the pUC19-transformed negative control to allow assessment of their depletion. For TATC, TGTC, TTCC, and TCCC PAMs, at least one AsCpf1 variant in the library was highly depleted ( $\geq 15$ -fold; **Fig. 1d** and **Supplementary Table 1b**). For TATC and TGTC, many of the depleted variants were at Lys548, a conserved residue that forms hydrogen bonds with the PAM duplex<sup>12,15</sup>. A number of hits were also observed for TTCC and TCCC, most notably an arginine substitution at Ser542, a non-conserved residue.

We evaluated whether variants identified in the screen had activity in HEK293T cells by targeting them to endogenous sites in two

genes (*DNMT1* and *VEGFA*; **Fig. 2a** and **Supplementary Fig. 1a**). Most of the variants we tested generated indels at target sites with their corresponding PAMs; in particular, K548V was most active at a TATC target site, whereas S542R markedly increased activity for two TTCC target sites as well as a TCCC site. Combining the top single amino acid mutations into double and triple mutants further improved activity (**Fig. 2a** and **Supplementary Fig. 1b**). We selected the variants with the highest activity, S542R/K607R (hereafter referred to as RR) and S542R/K548V/N552R (hereafter referred to as RVR), for further investigation.


To assess the global PAM preference of the RR and RVR variants and compare them with WT AsCpf1, we adapted an *in vitro* PAM identification assay described previously (**Fig. 2b**)<sup>1,16</sup>. We incubated cell lysate from HEK293T cells expressing AsCpf1 (or an engineered variant) with *in vitro*-transcribed crRNA and a library of plasmid DNA containing a constant target preceded by a degenerate sequence (5'-NNNNNNNNN-target). By amplifying and deep-sequencing the intact substrates and comparing them with the negative control, we determined which sequences were successfully cleaved. For each Cpf1 variant, nine reactions were carried out in parallel, each incubated for a different amount of time, in order to assess cleavage kinetics (**Supplementary Figs. 2** and **3**).

As expected, WT AsCpf1 was most active at TTTV PAMs (**Fig. 2c,d**) and had lower activity at TTTT, supporting the previously reported definition of the WT PAM as TTTV<sup>1,6</sup>. WT also cleaved other sequences including NTTV, TCTV, and TTCV at low rates, consistent with our observations in HEK293T cells (**Supplementary Fig. 4**) and in *E. coli*. By contrast, the RR and RVR variants had the highest activity at TYCV (where Y can be C or T) and TATV PAMs, respectively, compared to little or no activity for WT Cpf1 at those PAMs (**Fig. 2c,d**). The variant PAMs were also not as strictly defined as that of WT. The RR variant also cleaved ACCC and CCCC PAMs (and, to a lesser extent, VYCV), and the RVR variant also cleaved RATR PAMs (where R can be A or G).

To assess the robustness of the engineered PAM activity, we investigated the activity of the RR and RVR variants at their preferred PAMs (i.e., TYCV and TATV, respectively) across a diverse panel of endogenous target sites in HEK293T cells (**Fig. 2e** and **Supplementary Fig. 5**). The RR and RVR variants generated >50% indel for 14 out of 20 TYCV sites (70%) and 18 out of 23 TATV sites (78%), respectively, compared to little or no activity for WT AsCpf1 at most of these sites ( $P < 0.0001$  for both variants; Wilcoxon signed-rank). By comparison, WT AsCpf1 achieved >50% indel for 8 out of 23 TTTV sites (35%). These data suggest that, at their respective preferred PAMs, the variants have comparable or slightly higher activity than the WT nuclease (**Fig. 2e**). The RR variant also exhibited substantial rates of editing in mouse *Neuro2a* cells (>20% indel for 6 out of 9 TYCV sites) (**Supplementary Fig. 6**).

Based on our observations that the RR variant also cleaves VYCV PAMs *in vitro*, albeit at a substantially lower rate than TYCV, we tested the activity of the RR variant at a separate panel of VYCV sites in HEK293T cells (**Supplementary Fig. 4**). Across the four genes assessed (*CFTR*, *DNMT1*, *EMX1*, and *VEGFA*), the RR variant achieved >20% indel for 24 out of 36 VYCV sites (67%), suggesting that, when necessary, target sites with VYCV PAMs can also be considered for editing with the RR variant.

To quantify how these Cpf1 PAM variants affect the targeting range of the CRISPR-Cpf1 system, we performed a computational analysis of the distribution of PAM sequences in the human genome (**Fig. 2f** and **Supplementary Fig. 7**), excluding Ns and masked repeats. When considering only the most active PAMs, the variants and WT collectively expand the targeting range of Cpf1 to one target site per ~11 bp in human coding sequences (corresponding to an approximately threefold increase relative to WT alone) and reduce the median distance to the



**FIGURE 2** Construction and characterization of AsCpf1 variants with altered PAM specificities. (a) Combinatorial mutagenesis identifies AsCpf1 variants that cleave target sites with TYCV and TATV PAMs in HEK293T cells, where Y = C or T, and V = A, C, or G (see also **Supplementary Fig. 1**). Error bars: mean  $\pm$  s.e.m. for  $n = 4$  transfected cell cultures. (b) Schematic of *in vitro* cleavage assay used to determine global PAM specificity (see also **Supplementary Figs. 2 and 3**). (c) Web logos of the most rapidly cleaved PAMs for wild-type (WT), S542R/K607R (RR), and S542R/K548V/N552R (RVR) variants. (d) Normalized cleavage rates for all 4-base PAMs for WT and variants. NNRN PAMs are not shown due to negligible cleavage. The most active PAMs are boxed in red. (e) Comparison of the activity of WT, RR, and RVR at their preferred PAMs at a diverse panel of target sites in HEK293T cells (see also **Supplementary Fig. 5**). Each dot represents a target site. For indel percentages, each dot represents the mean of  $n = 3$  transfected cell cultures, and the red lines indicate the overall means within each group. For fold change, each dot represents the ratio of the means of the corresponding indel replicates. n.s.  $P > 0.05$  (Mann–Whitney); \* $P < 0.05$  (Mann–Whitney); \*\*\*\* $P < 0.0001$  (Wilcoxon signed-rank). (f) Targeting range of AsCpf1 variants in the human genome and in coding sequences (see also **Supplementary Fig. 7**). Plots show the probability mass function of the distance in base pairs to the nearest cleavage site. The boxplots indicate median and interquartile range. Genomic regions that contain Ns or masked repeats were ignored.

nearest cleavage site to 3 bp. Moreover, when considering a more broadly defined set of efficiently cleavable PAMs (in particular, the preferred PAMs plus MCCC and RATR, where M can be A or C), the targeting range is further expanded to one site per  $\sim 7$  bp in human coding sequences, with a median distance to the nearest cleavage site of 2 bp.

We evaluated the genome-wide editing specificity of the RR and RVR variants using BLISS (double-strand breaks labeling *in situ* and sequencing), which quantifies DNA double-stranded breaks (DSBs) across the genome<sup>7</sup>. To fairly compare the variants to WT, we selected target sites bearing PAMs that can be reliably cleaved by all three nucleases; TTTV was the only PAM that met this criterion, although it has lower activity for the RR variant. For three of the four target sites evaluated (*VEGFA*, *GRIN2B*, and *DNMT1*), no off-target activity was detected from deep sequencing of the BLISS-identified loci (**Fig. 3a** and **Supplementary Table 2**) for any of the nucleases. For the fourth target site (*EMX1*), BLISS identified six off-target sites with detectable indels; all six sites had a TTCA PAM and no more than one mismatch in the first 19 nucleotides of the guide. As expected, both variants had increased activity at these off-target sites compared to WT, consistent with their increased ability to recognize TTCA PAMs. On the other hand, when targeting a site in the *RPL32P3* gene with known TTTV off-target sites<sup>5</sup>, the variants exhibited similar or reduced off-target activity (**Fig. 3b**), which is also consistent with PAM preference. Collectively, these results indicate that the variants retain a high level of editing specificity that is comparable to WT AsCpf1. We note that a few of the off-target sites with low indel frequencies were not detected by BLISS at the time point we sampled, likely reflecting the dependence of BLISS on the timing of DSB formation<sup>7</sup>.


We investigated whether the specificity of AsCpf1 can be improved by removing non-specific contacts between positively charged or

polar residues and the target DNA, similar to strategies previously employed with *Streptococcus pyogenes* Cas9 (SpCas9)<sup>17,18</sup>. We identified K949A, which is located in the cleft of the protein that is hypothesized to interact with the non-target DNA strand, as a candidate (**Supplementary Fig. 8**). When combined with the RR and RVR variants, K949A reduced cleavage at all off-target sites assessed (**Fig. 3c**) while maintaining high levels of on-target activity (**Fig. 3d**).

Because Cpf1-family endonucleases have strong sequence and structural homology, the 542, 548, 552, and 607 positions in AsCpf1 have clear correspondences in other Cpf1 orthologs (**Supplementary Fig. 9** and **Supplementary Table 3**). Based on sequence alignment and the crystal structure<sup>19</sup>, we hypothesized that LbCpf1 could also be engineered to recognize TYCV and TATV PAMs by introducing the mutations G532R/K595R and G532R/K538V/Y542R, respectively (**Supplementary Fig. 10a**). These mutations altered the PAM specificity of LbCpf1 in the predicted manner (**Supplementary Figs. 10b and 11**), suggesting that this approach may be generally applicable across Cpf1 orthologs.

In summary, we have demonstrated that despite its evolutionary conservation, the PAM preference of Cpf1-family endonucleases can be altered by suitable mutations to residues close to the PAM duplex. Using a structure-guided mutagenesis screen, we engineered two variants, RR and RVR, which can robustly cleave target sites with TYCV and TATV PAMs, respectively, in mammalian cells. We extended this approach to similarly modify a second Cpf1 ortholog. Finally, we introduced an additional mutation that enhanced Cpf1 specificity. Collectively, these engineered variants increase the targeting range of Cpf1 to one cleavage site for every  $\sim 11$  bp in human coding sequences and provide useful additions to the CRISPR–Cas genome engineering toolbox.





**Figure 3** Specificity of AsCpf1 PAM variants. (a) DNA double-strand breaks labeling *in situ* and sequencing (BLISS) for four target sites (*VEGFA*, *GRIN2B*, *EMX1*, and *DNMT1*) in HEK293T cells. The  $\log_{10}$  number of unique double-strand break (DSB) ends per  $10^5$  reads is indicated by the magenta heat map. The normalized PAM cleavage rates from the *in vitro* cleavage assay in **Figure 2d** are indicated by the blue heat map. Each BLISS-identified cleavage site was independently assessed for indel formation (bar graphs). Bars show mean  $\pm$  s.e.m. for  $n = 4$  transfected cell cultures. Mismatches in bases 21–23 of the target are grayed as they have minimal impact on cleavage efficiency<sup>4,5</sup>. (b) Evaluation of an additional target site in the *RPL32P3* gene with known TTTV off-target sites<sup>5</sup>. (c) Addition of a K949A mutation improves the specificity of WT AsCpf1 and variants (see also **Supplementary Fig. 8**). For **b** and **c**, bars show mean  $\pm$  s.e.m. for  $n = 3$  transfected cell cultures. (d) On-target efficiency of the RR and RVR variants  $\pm$  K949A. Each dot represents a distinct target site (mean of  $n = 3$  transfected cell cultures).

## METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the [online version of the paper](#).

Note: Any Supplementary Information and Source Data files are available in the [online version of the paper](#).

## ACKNOWLEDGMENTS

We thank A. Magnell for experimental assistance; R. Macrae for a critical review of the manuscript; and the entire Zhang laboratory for support and advice. D.B.T.C. is supported by T32GM007753 from the National Institute of General Medical Sciences. W.X.Y. is supported by T32GM007753 from the National Institute of General Medical Sciences and a Paul and Daisy Soros Fellowship. J.C.M. is supported by the NIH (training grant 2 T32 GM 7287-41). H.N. is supported by JST, PRESTO (JPMJPR13L8), JSPS KAKENHI (Grant Numbers 26291010 and 15H01463). O.N. is supported by the Basic Science and Platform Technology Program for Innovative Biological Medicine from the Japan Agency for Medical Research and Development, AMED, and the Council for Science, and Platform for Drug Discovery, Informatics, and Structural Life Science from the Ministry of Education, Culture, Sports, Science and Technology. N.C. is supported by the Karolinska Institutet, the Swedish Research Council (521-2014-2866), the Swedish Cancer Research Foundation (CAN 2015/585), and the Ragnar Söderberg Foundation. E.Z. is a New York Stem Cell Foundation–Robertson Investigator. E.Z. is supported by the NIH through NIMH (5DP1-MH100706 and 1R01-MH110049), NSF, Howard Hughes Medical Institute, the New York Stem Cell, Simons, Paul G. Allen Family, and Vallee Foundations; and James and Patricia Poitras, Robert Metcalfe, and David Cheng.

## AUTHOR CONTRIBUTIONS

L.G., D.B.T.C., and E.Z. conceived this study. L.G. and D.B.T.C. performed experiments with help from all authors. J.C.M. contributed to the bacterial selection screen. M.W.S. processed BLISS samples, and W.X.Y. analyzed BLISS data. T.Y., H.N., and O.N. provided unpublished AsCpf1 crystal structure information. N.C. provided an unpublished BLISS protocol. E.Z. supervised research. L.G. and E.Z. wrote the manuscript with input from all authors.

## COMPETING FINANCIAL INTERESTS

The authors declare competing financial interests: details are available in the [online version of the paper](#).

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

- Zetsche, B. *et al.* Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR–Cas system. *Cell* **163**, 759–771 (2015).
- Fonfara, I., Richter, H., Bratovič, M., Le Rhun, A. & Charpentier, E. The CRISPR-associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA. *Nature* **532**, 517–521 (2016).
- Zetsche, B. *et al.* Multiplex gene editing by CRISPR–Cpf1 using a single crRNA array. *Nat. Biotechnol.* **35**, 31–34 (2017).
- Kim, D. *et al.* Genome-wide analysis reveals specificities of Cpf1 endonucleases in human cells. *Nat. Biotechnol.* **34**, 863–868 (2016).
- Kleinstiver, B.P. *et al.* Genome-wide specificities of CRISPR–Cas Cpf1 nucleases in human cells. *Nat. Biotechnol.* **34**, 869–874 (2016).
- Kim, H.K. *et al.* *In vivo* high-throughput profiling of CRISPR–Cpf1 activity. *Nat. Methods* **14**, 153–159 (2017).
- Yan, W.X. *et al.* BLISS is a versatile and quantitative method for genome-wide profiling of DNA double-strand breaks. *Nat. Commun.* **8**, 15058 (2017).
- Kleinstiver, B.P. *et al.* Engineered CRISPR–Cas9 nucleases with altered PAM specificities. *Nature* **523**, 481–485 (2015).
- Kleinstiver, B.P. *et al.* Broadening the targeting range of *Staphylococcus aureus* CRISPR–Cas9 by modifying PAM recognition. *Nat. Biotechnol.* **33**, 1293–1298 (2015).
- Hirano, S., Nishimasu, H., Ishitani, R. & Nureki, O. Structural basis for the altered PAM specificities of engineered CRISPR–Cas9. *Mol. Cell* **61**, 886–894 (2016).
- Anders, C., Bargsten, K. & Jinek, M. Structural plasticity of PAM recognition by engineered variants of the RNA-guided endonuclease Cas9. *Mol. Cell* **61**, 895–902 (2016).
- Yamano, T. *et al.* Crystal structure of Cpf1 in complex with guide RNA and target DNA. *Cell* **165**, 949–962 (2016).
- Jiang, W., Bikard, D., Cox, D., Zhang, F. & Marraffini, L.A. RNA-guided editing of bacterial genomes using CRISPR–Cas systems. *Nat. Biotechnol.* **31**, 233–239 (2013).
- Esvelt, K.M. *et al.* Orthogonal Cas9 proteins for RNA-guided gene regulation and editing. *Nat. Methods* **10**, 1116–1121 (2013).
- Gao, P., Yang, H., Rajashankar, K.R., Huang, Z. & Patel, D.J. Type V CRISPR–Cas Cpf1 endonuclease employs a unique mechanism for crRNA-mediated target DNA recognition. *Cell Res.* **26**, 901–913 (2016).
- Ran, F.A. *et al.* *In vivo* genome editing using *Staphylococcus aureus* Cas9. *Nature* **520**, 186–191 (2015).
- Slaymaker, I.M. *et al.* Rationally engineered Cas9 nucleases with improved specificity. *Science* **351**, 84–88 (2016).
- Kleinstiver, B.P. *et al.* High-fidelity CRISPR–Cas9 nucleases with no detectable genome-wide off-target effects. *Nature* **529**, 490–495 (2016).
- Dong, D. *et al.* The crystal structure of Cpf1 in complex with CRISPR RNA. *Nature* **532**, 522–526 (2016).

## ONLINE METHODS

**Library construction.** Human codon-optimized AsCpf1 driven by a T7 promoter was cloned into a modified pACYC backbone, and unique restriction sites were introduced flanking the selected PAM-proximal AsCpf1 residues via suitable silent mutations. For each residue, a mutagenic insert was synthesized as short complementary oligonucleotides (Integrated DNA Technologies), with the mutated codon replaced by a degenerate NNK mixture of bases (where K can be G or T). Each degenerate codon position was also barcoded by creating a unique combination of silent mutations in neighboring codons in order to correct for sequencing errors during screen readout. The variant library was assembled by cassette mutagenesis, mini-prepped, pooled, and precipitated with isopropanol.

***E. coli* negative-selection screen.** NovaBlue(DE3) *E. coli* (Novagen) cells were transformed with the variant library and plated on LB agar (Affymetrix) containing 25 µg/mL chloramphenicol. Surviving colonies were scraped and cultured in ZymoBroth with 25 µg/mL chloramphenicol to an O.D. of 0.4–0.6 and made competent using a Mix & Go kit (Zymo). For each mutant PAM screened, the competent *E. coli* pool was transformed with 100-ng target plasmid containing the mutant PAM, incubated on ice for 15–30 min, heat shocked at 42 °C for 30s, and plated on LB agar containing 100 µg/mL ampicillin and 25 µg/mL chloramphenicol in the absence of IPTG. A negative control was obtained by transforming the *E. coli* with pUC19, which lacks the target site. Plasmid DNA from surviving colonies was isolated by midi-prep (Qiagen). The regions containing mutations were amplified with custom primers containing Illumina adaptors and paired-end sequenced with a 600-cycle MiSeq kit (Illumina). Reads were filtered by requiring perfect matches to silent codon barcodes; a Phred quality (Q score) of at least 30 for each of the three NNK bases; and consistency between forward and reverse reads, when applicable. The read count for each variant was normalized assuming that the mean abundance of TAG (stop) codons was equivalent to the negative control.

***In vitro* PAM identification assay.** Plasmids encoding the AsCpf1 variants were transfected into HEK293T cells as described below. Cell lysate was prepared with lysis buffer (20 mM HEPES, 100 mM KCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 5% glycerol, 0.1% Triton X-100) supplemented with EDTA-free cOmplete Protease Inhibitor Cocktail (Roche). crRNA was transcribed *in vitro* using custom oligonucleotides and HiScribe T7 *in vitro* Transcription Kit (NEB) following the manufacturer's recommended protocol. For the PAM library, a degenerate 8-bp sequence preceding a 33-bp target site<sup>1</sup> was cloned into the multiple cloning site in pUC19, and the library was digested with AatII and LguI and gel purified before use. Each *in vitro* cleavage reaction consisted of 1 µL 10× CutSmart buffer (NEB), 25 ng PAM library, 250 ng *in vitro*-transcribed crRNA, 0.5 µL cell lysate, and water for a total volume of 10 µL. Reactions were incubated at 37 °C and quenched by adding 50 µL Buffer PB (Qiagen) followed by column purification. Purified DNA was amplified with two rounds of PCR over 29 total cycles using custom primers containing Illumina adaptors and sequenced with a 75-cycle NextSeq kit (Illumina). For each Cpf1 variant, separate *in vitro* cleavage reactions were carried out for 1.15 min, 4 min, 10 min, 15 min, 20 min, 30 min, 40 min, 90 min, and 175 min. The unmodified library of degenerate sequences was used as the 0 min time point. A negative control, using lysate from unmodified HEK293T cells, was taken at 10 min.

**Computational analysis of PAM cleavage kinetics.** See also **Supplementary Figures 2 and 3**. Sequencing reads were filtered by Phred quality ( $\geq 30$  for all of the eight degenerate PAM bases). For each cleavage reaction, a depletion ratio for each of the 4<sup>8</sup> PAM sequences was calculated as (normalized read count in cleavage reaction)/(normalized read count in negative control). Each depletion ratio was then divided by the median depletion ratio of all NNNNVRTT sequences, which were not cleaved by WT AsCpf1 or either of the variants. The depletion ratios of each PAM sequence (4<sup>8</sup> total) across time points were fit using nonlinear least-squares to an exponential decay model  $x(t) = c_0 + ce^{-kt}$ , where  $x(t)$  is the depletion ratio at time  $t$ , and the terms  $c_0 \leq 0.2$ ,  $c$ , and  $k$  (the rate constant in min<sup>-1</sup>) are parameters. For each variant, the estimated cleavage

rate  $k$  of each 4-base PAM was computed as the median cleavage rate of the 256 8-base sequences corresponding to that PAM; for instance, the cleavage rate of TTTA was computed as the median cleavage rate of the 256 sequences of the form NNNNTTTA. Finally, all cleavage rates were adjusted so that the highest rate of any 4-base PAM was equal to 1 for each variant.

**Cell culture and transfection.** Human embryonic kidney 293 and Neuro2a cell lines were maintained in Dulbecco's modified Eagle's medium supplemented with 10% FBS (Gibco) at 37 °C with 5% CO<sub>2</sub> incubation. Cells were seeded one day before transfection in 24- or 96-well plates (Corning) at a density of approximately  $1.2 \times 10^5$  cells per 24-well or  $2.4 \times 10^4$  cells per 96-well and transfected at 50–80% confluency using Lipofectamine 2000 (Invitrogen), according to the manufacturer's recommended protocol. For cell lysates, 500 ng of Cpf1 plasmid was delivered per 24-well. For indel analysis in HEK293T cells, 400 ng of Cpf1 plasmid plus 100 ng crRNA plasmid was delivered per 24-well, or 100 ng Cpf1 plus 50 ng crRNA plasmid per 96-well. For BLISS and for indel analysis in Neuro2a cells, 500 ng of a plasmid encoding both Cpf1 and crRNA were delivered per 24-well. All indel and BLISS experiments used a guide length of 23 nucleotides.

**Indel quantification.** All indel frequencies were quantified by targeted deep sequencing (Illumina). For indel library preparation, cells were harvested approximately 3 days after transfection, and genomic DNA was extracted using QuickExtract DNA extraction solution (Epicentre) by resuspending pelleted cells in QuickExtract (80 µL per 24-well, or 20 µL per 96-well), followed by incubation at 65 °C for 15 min, 68 °C for 15 min, and 98 °C for 10 min. Amplicons for deep sequencing were generated using two rounds of PCR to attach Illumina handles. Indels were counted computationally by searching each sequencing read for exact matches with strings delineating the ends of a 50- to 70-bp window around the cut site. The distance in bp between these strings was then compared to the corresponding distance in the reference genome, and the read was counted as an indel if the two distances differed. For each sample, the indel frequency was determined as (number of reads with an indel)/(number of total reads). Samples with fewer than 1,000 total reads were excluded. Where negative control data are not shown, indel percentages represent background-subtracted maximum likelihood estimates. In particular, for a sample with  $R$  total reads, of which  $n \leq R$  are indels, and false-positive rate  $0 \leq \alpha < 1$  (as determined by the negative control), the true indel rate was estimated as  $\max\{0, [(n/R) - \alpha]/(1 - \alpha)\}$ .

**Computational analysis of Cpf1 targeting range.** The complete GRCh38 human genome assembly and coding sequences, with repeats and low complexity regions masked, were downloaded from Ensembl and analyzed as described in **Supplementary Figure 7**.

**BLISS.** All BLISS experiments and analyses were performed as previously described<sup>7</sup>. The data analysis for the staggered cut sites of Cpf1 was slightly modified from prior analysis<sup>7,16</sup> to increase sensitivity. Previously, to distinguish bona fide nuclease-induced events from the background DSBs in DSB hotspots, centromeres, and telomeres, we had used a cutoff based on the fraction of the pairwise reads that overlapped less than -6 bp. This cutoff was set at 0.95 based on empirical data from Cas9 off-target analysis, but to accommodate the variation produced by the staggered cut sites of Cpf1, we found that greater sensitivity to Cpf1 off-target sites could be obtained by relaxing this cutoff to 0.85. All other analyses, such as the guide homology score calculations, were as described<sup>7</sup>.

**Plasmids and guide sequences.** A list of the plasmids and guide sequences used in this study can be found in **Supplementary Table 4**.

**Data availability.** Deep sequencing data are available on the NCBI Sequence Read Archive [SRR5611789](https://www.ncbi.nlm.nih.gov/sra/SRR5611789). Reagents and further information will be available to the academic community through Addgene and the Zhang laboratory website (<http://www.genome-engineering.org/>).